

**RESEARCH AND MANAGEMENT OF PACIFIC RIDLEY SEATURTLES,
(*Lepidochelys olivacea*) -- MARINE TURTLE RESEARCH CENTRE, PURI, ORISSA**
Sudhakar Kar, Research Officer, C/o CCF (WL), Forest Department of Orissa, Bhubaneswar-1

INTRODUCTION

Following the advice of Dr. H. R. Bustard (FAO, 1975), the Forest Department of Orissa became actively involved in conservation, research and management of endangered Pacific Ridley Sea Turtles (*Lepidochelys olivacea*) at Gahirmatha coast of Bhitarkanika Wildlife Sanctuary during the year 1976.

The aims of the Project were two-fold (Kanungo, B.C., Forest Deptt., Orissa, 1976):

1.(a) To conserve this endangered reptilian species by providing adequate protection to the sea turtles both on the shore and at off-shore areas, and (b) to conduct research on the ecology and biology of sea turtles.

2. Later, to exploit the species for earning the much needed foreign exchange.

Since then, sincere attempts have been made to accumulate more scientific data on nest/nesting, egg laying, incubation of eggs, hatching success, predation of eggs and hatchlings, tagging of nesting female turtle, tag recovery and other associated ecological studies on Pacific Ridley Sea turtles at Gahirmatha beach.

Due to several adverse factors such as location of the place, communication difficulty, the want of electricity facility, soil condition, tidal bore, etc. it became difficult to conduct a few important studies on turtle's biology at Gahirmatha. So an alternate site for the establishment of a research station was selected and the Seaturtle Research Project was broadly divided into two components :

1. In situ research on the ecology of Pacific Ridley Turtles at Gahirmatha rookery of Bhitarkanika Wildlife Sanctuary. (It is the largest sea turtle rookery of its kind in the world (Bustard, 1976). A minimum of 2.0 lakhs to a maximum of 6.5 lakhs with an average of 2.5 lakhs of nesting female turtles visit the beach every year for nesting and egg laying).

2. Off-shore research on the biology of the Pacific Ridley turtles.

OBJECTIVES

This off-shore Research Centre was established at Beleswarpatana on the bank of the river Nuanai, 10 km. away from Puri along the Puri-Konark marine-drive road within the Balukhand Wildlife Sanctuary during the year 1985 - 86.

The basic objectives of this Sea Turtle Research Centre are to study on the following important aspects:

- i. Environmental Sex Determination (ESD) of sea turtles by incubating the eggs at controlled temperature and moisture conditions. (This is most important aspect of study for future management of sea turtles).
- ii. Collection of eggs from "sporadic" nests from adjacent beaches, incubation of eggs under simulated conditions and study on incubation and hatching success, etc.
- iii. Food and feeding habits and food preference in captivity.
- iv. Food vis-a-vis growth of turtles in captivity to highlight food conversion.
- v. Behaviour of the turtles of various year classes with change of seasons.
- vi. Health problems and preventive measures in captive husbandry conditions.
- vii. Seasonal change of salinity in the river/sea water and its impact in the physiology of turtles.
- viii. Age of maturity, and captive breeding of sea turtles which is so far an intriguing aspect of study to the biologists.
- ix. To provide educative and recreational facilities to the public and tourists.

ACHIEVEMENTS

The Marine Turtle Research Centre was started with only 26 turtles of about two years old (1984 hatching), which were brought from Gahirmatha Marine Turtle Research Centre on 26.1.86. Those turtles are accommodated in plastic pools of different diameters and brackish water from the nearby Nuanai river is pumped into the pools on alternate days. Proper hygienic conditions are maintained in and around the pools and the pool water is being kept clean. Turtles are fed daily with marine fish and prawns (Table 1).

At the beginning there was casualty among the turtles. Several causes are being attributed for their sickness and death, the major being the want of clear saline water (clear sea water) in the rearing pools. The river water is remaining dirty/turbid during the entire rainy season (June to October) and salinity of the river water decreases to degree zero in the months of July and September. After rains, the salinity gradually increases (November onwards) and reaches the maximum in hot summer months (April and May).